

PicoAg–BFIV-25B-Arizona

Bactericide, Fungicide, Viricide, Insecticide For Agriculture Use

Active Ingredient: Zinc 1%.

Inert Ingredients: Carbon 94%, Nitrogen 5%.

KEEP OUT OF REACH OF CHILDREN

FIFRA Exempt Statement Preferred Statements. This product has not been registered by the United States Environmental Protection Agency. [Biobased USA] represents that his product qualifies for exemption from registration under FIFRA. This product qualifies for exemption from EPA registration under Federal Insecticide, Fungicide and Rodenticide Act (FIFRA)”

Precautionary Statements, First Aid, and other Hazard Statements Restrictions: Personal Protective Equipment not required, this product is non-hazardous. Physician and Fire Department. Long term, widespread usage and review of ingredients indicate this product to be generally regarded as a safe (GRAS) product. There is no specific antidote. Treatment of overexposure should be directed at the control of symptoms and the clinical condition. No Fire and or exposure hazards.

Directions for Use: For vegetables apply 2.5 ounces in 10 gallons per acre. Repeat every 20 days but skip application while plant flowering.

Direction of Application for major crops: Apply 5 oz to 8 oz per 10 gallons of water per acre. Timing for application is at Seed Planting or 3 leaf, 30 and 60 days after planting stages of plant growth for best results. Consult with our technical staff for crop specific application instructions. Aerial Spraying is 5 oz to 5 gallons.

Pests Controlled:

Bacteria: Acetobacter, Acinetobacter, Alfalfa Wilt, Actinomyces, Agrobacterium, Alcaligenes, Bordetella, Candida, Tomato Bacterial Canker, Cardiobacterium, Chlamydia, Clostridium, Corynebacterium, Coxiella, Diplococcus, Erwinia, E Coli, Francisella, Gummosis, Goss’s Wilt (Leaf blight), Haemophilus, Klebsiella, Listeria, Methicillin Resistant Staphylococcus Aureus (MRSA), Micromonospora, Moko Disease, Moraxella, Mycobacterium, Mycoplasma, Neisseria, Nocardia, Pediococcus, Pneumocystis, Pseudomonas, Rice blight, Salmonella, Sarcina, Serratia, Shigella, Staphylococcus, Streptococcus, Thermoactinomyces, Thermomonospora, Xanthomonas. Clavibacter michiganensis subsp. californiensis Clavibacter michiganensis subsp. capsici Clavibacter michiganensis subsp. chilensis Clavibacter michiganensis subsp. insidiosus Clavibacter michiganensis subsp. michiganensis Clavibacter michiganensis subsp. nebraskensis Clavibacter michiganensis subsp. phaseoli Clavibacter michiganensis subsp. sepedonicus Clavibacter michiganensis subsp. tessellarius

Fungi: Absidia, Acremonium, Alternaria, Antennariella, Anthracnose, Ascomycete, Aspergillus, Atheliaceae, Aureobasidium, Black leaf streak, Black Shank, Black Sigatoka, Black Spot, Blastomyces, Botryosphaeriaceae, Botrytis, Capnodium, Cercospora,

Chaetomium, Cladosporium, Coccidioides, Cochliobolus, Coffee Rust, Common Smut, Corn Blight, Corn Leaf Blight, Corn Leaf Spot, Cryptococcus, Diplocarpon, Dracaena, Emericella, Enterobacteriaceae, Epicoccum, Erysiphales, Eurotium, Exophiala, Eyespot, Fusarium Geomyces, Gray Leaf Spot, Helminthosporium, Histoplasma, Leaf Blight, Leaf Spot, Limacinula, Macrophomina, Mango Malformation, Meliola, Mucor plumbeus, Mycena Citricolor, Mycosphaerella, Necrosis, Rust, Oomycete, Paecilomyces, Panama Disease, Paracoccidioides Penicillium, Phialophora, Phoma, Phomopsis, Phytophthora, Podosphaera, Powdery Mildew, Puccinia, Rhizoctonia, Rhodoturula, Rice blast, Rice sheath, Rust, Sclerotinia, Scopulariopsis, Scorias, Sooty mold, Sporothrix, Stachybotrys, Stem Rust, Stripe rust, Sudden Death Syndrome (SDS), Trichoderma, Ulocadium, Volutella, Wallemia, White mold, Witches' Broom, Yellow Sigatoka.

Viruses: Abacá bract mosaic virus, Abaca bunchy top virus, Abaca mosaic virus, Adenovirus Virus, Banana mild mosaic virus, Banana Streak Virus, Bean Pod Mottle Virus, Bract mosai virus, Banana bunchy top virus, Chlamydia pneumoiae Virus, Coxsackievirus Virus, Cucumber Mosaic Virus, Echovirus Virus, Hantavirus Virus, Morbillivirus Virus, Papaya Ring Spot Virus, Parainfluenza Virus, Paramyxovirus Virus, Parovirus B19 Virus, Poxvirus, Rhinovirus Virus, Sharka Plum Pox Virus, Soybean Mosaic Virus, Togavirus Virus, Tomato Leaf Curl, Tomato yellow leaf curl virus, Varicella-zoster Virus, Yersinia pestis Virus.

Insects: Aphids, Army worms, Cereal Leaf Beetle, Japanese Beetle, Leaf Hopper, Leaf Miner, Mealy Bugs, Red Mites, Scale Euonymus, Scale False Oleander, Scale Green, Scale Red, Scale San Jose, Scale Soft Green, Scale Torpedo, Spider mites Spotted Mites, Stink Bugs, Thrips, White Flies, Wooly Adelgid.

This product is not required to be registered by the United States Environmental Protection Agency. Biobased USA represents that this product qualifies for exemption from registration under FIFRA.

Storage: Store in a cool, well-ventilated area.

Disposal: Evacuate the spill area as floor may be slippery. Use care to prevent falling. Flush area with water. Dispose of as per local, state and federal regulations.

Health Hazard Acute Exposure. No evidence of adverse effects from: Swallowing, Skin Absorption, Skin Contact, Eye Contact, Inhalation.

Transportation: Non Dangerous Goods Not Regulated, Non Hazardous.

Net Contents: Packing can be 1 gallon, 8 lbs., 2.5 gallons, 20 pounds, 30 gallon drum, 240 pounds, 275 gallon 1040 liter totes.

Biobased USA, 1611 Speer Bridge Road, Yadkinville, NC 27055
Donald R. Wilshe, CEO. [Email: donwilshe@biobased.us](mailto:donwilshe@biobased.us)
Phone(24/7): 800 995-9203, +1 (336) 306-0193